



## **AP® Calculus BC 2009 Free-Response Questions**

### **The College Board**

The College Board is a not-for-profit membership association whose mission is to connect students to college success and opportunity. Founded in 1900, the association is composed of more than 5,600 schools, colleges, universities and other educational organizations. Each year, the College Board serves seven million students and their parents, 23,000 high schools and 3,800 colleges through major programs and services in college readiness, college admissions, guidance, assessment, financial aid, enrollment, and teaching and learning. Among its best-known programs are the SAT®, the PSAT/NMSQT® and the Advanced Placement Program® (AP®). The College Board is committed to the principles of excellence and equity, and that commitment is embodied in all of its programs, services, activities and concerns.

© 2009 The College Board. All rights reserved. College Board, Advanced Placement Program, AP, AP Central, SAT, and the acorn logo are registered trademarks of the College Board. PSAT/NMSQT is a registered trademark of the College Board and National Merit Scholarship Corporation.


Permission to use copyrighted College Board materials may be requested online at:  
[www.collegeboard.com/inquiry/cbpermit.html](http://www.collegeboard.com/inquiry/cbpermit.html).

Visit the College Board on the Web: [www.collegeboard.com](http://www.collegeboard.com).  
AP Central is the official online home for the AP Program: [apcentral.collegeboard.com](http://apcentral.collegeboard.com).

**2009 AP® CALCULUS BC FREE-RESPONSE QUESTIONS**

**CALCULUS BC  
SECTION II, Part A  
Time—45 minutes  
Number of problems—3**

**A graphing calculator is required for some problems or parts of problems.**



1. Caren rides her bicycle along a straight road from home to school, starting at home at time  $t = 0$  minutes and arriving at school at time  $t = 12$  minutes. During the time interval  $0 \leq t \leq 12$  minutes, her velocity  $v(t)$ , in miles per minute, is modeled by the piecewise-linear function whose graph is shown above.
  - (a) Find the acceleration of Caren's bicycle at time  $t = 7.5$  minutes. Indicate units of measure.
  - (b) Using correct units, explain the meaning of  $\int_0^{12} |v(t)| dt$  in terms of Caren's trip. Find the value of  $\int_0^{12} |v(t)| dt$ .
  - (c) Shortly after leaving home, Caren realizes she left her calculus homework at home, and she returns to get it. At what time does she turn around to go back home? Give a reason for your answer.
  - (d) Larry also rides his bicycle along a straight road from home to school in 12 minutes. His velocity is modeled by the function  $w$  given by  $w(t) = \frac{\pi}{15} \sin\left(\frac{\pi}{12}t\right)$ , where  $w(t)$  is in miles per minute for  $0 \leq t \leq 12$  minutes. Who lives closer to school: Caren or Larry? Show the work that leads to your answer.

---


**WRITE ALL WORK IN THE PINK EXAM BOOKLET.**

## **2009 AP<sup>®</sup> CALCULUS BC FREE-RESPONSE QUESTIONS**

2. The rate at which people enter an auditorium for a rock concert is modeled by the function  $R$  given by  $R(t) = 1380t^2 - 675t^3$  for  $0 \leq t \leq 2$  hours;  $R(t)$  is measured in people per hour. No one is in the auditorium at time  $t = 0$ , when the doors open. The doors close and the concert begins at time  $t = 2$ .
- (a) How many people are in the auditorium when the concert begins?
- (b) Find the time when the rate at which people enter the auditorium is a maximum. Justify your answer.
- (c) The total wait time for all the people in the auditorium is found by adding the time each person waits, starting at the time the person enters the auditorium and ending when the concert begins. The function  $w$  models the total wait time for all the people who enter the auditorium before time  $t$ . The derivative of  $w$  is given by  $w'(t) = (2 - t)R(t)$ . Find  $w(2) - w(1)$ , the total wait time for those who enter the auditorium after time  $t = 1$ .
- (d) On average, how long does a person wait in the auditorium for the concert to begin? Consider all people who enter the auditorium after the doors open, and use the model for total wait time from part (c).
- 

**WRITE ALL WORK IN THE PINK EXAM BOOKLET.**

**2009 AP<sup>®</sup> CALCULUS BC FREE-RESPONSE QUESTIONS**



Note: Figure not drawn to scale.

3. A diver leaps from the edge of a diving platform into a pool below. The figure above shows the initial position of the diver and her position at a later time. At time  $t$  seconds after she leaps, the horizontal distance from the front edge of the platform to the diver's shoulders is given by  $x(t)$ , and the vertical distance from the water surface to her shoulders is given by  $y(t)$ , where  $x(t)$  and  $y(t)$  are measured in meters. Suppose that the diver's shoulders are 11.4 meters above the water when she makes her leap and that

$$\frac{dx}{dt} = 0.8 \text{ and } \frac{dy}{dt} = 3.6 - 9.8t,$$

for  $0 \leq t \leq A$ , where  $A$  is the time that the diver's shoulders enter the water.

- (a) Find the maximum vertical distance from the water surface to the diver's shoulders.
- (b) Find  $A$ , the time that the diver's shoulders enter the water.
- (c) Find the total distance traveled by the diver's shoulders from the time she leaps from the platform until the time her shoulders enter the water.
- (d) Find the angle  $\theta$ ,  $0 < \theta < \frac{\pi}{2}$ , between the path of the diver and the water at the instant the diver's shoulders enter the water.

**WRITE ALL WORK IN THE PINK EXAM BOOKLET.**

**END OF PART A OF SECTION II**

**2009 AP® CALCULUS BC FREE-RESPONSE QUESTIONS**

**CALCULUS BC  
SECTION II, Part B  
Time—45 minutes  
Number of problems—3**

**No calculator is allowed for these problems.**

4. Consider the differential equation  $\frac{dy}{dx} = 6x^2 - x^2y$ . Let  $y = f(x)$  be a particular solution to this differential equation with the initial condition  $f(-1) = 2$ .
- (a) Use Euler's method with two steps of equal size, starting at  $x = -1$ , to approximate  $f(0)$ . Show the work that leads to your answer.
- (b) At the point  $(-1, 2)$ , the value of  $\frac{d^2y}{dx^2}$  is  $-12$ . Find the second-degree Taylor polynomial for  $f$  about  $x = -1$ .
- (c) Find the particular solution  $y = f(x)$  to the given differential equation with the initial condition  $f(-1) = 2$ .
- 

**WRITE ALL WORK IN THE PINK EXAM BOOKLET.**

**2009 AP® CALCULUS BC FREE-RESPONSE QUESTIONS**

$x$	2	3	5	8	13
$f(x)$	1	4	-2	3	6

5. Let  $f$  be a function that is twice differentiable for all real numbers. The table above gives values of  $f$  for selected points in the closed interval  $2 \leq x \leq 13$ .
- (a) Estimate  $f'(4)$ . Show the work that leads to your answer.
- (b) Evaluate  $\int_2^{13} (3 - 5f'(x)) dx$ . Show the work that leads to your answer.
- (c) Use a left Riemann sum with subintervals indicated by the data in the table to approximate  $\int_2^{13} f(x) dx$ . Show the work that leads to your answer.
- (d) Suppose  $f'(5) = 3$  and  $f''(x) < 0$  for all  $x$  in the closed interval  $5 \leq x \leq 8$ . Use the line tangent to the graph of  $f$  at  $x = 5$  to show that  $f(7) \leq 4$ . Use the secant line for the graph of  $f$  on  $5 \leq x \leq 8$  to show that  $f(7) \geq \frac{4}{3}$ .
- 

6. The Maclaurin series for  $e^x$  is  $e^x = 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \cdots + \frac{x^n}{n!} + \cdots$ . The continuous function  $f$  is defined by

$$f(x) = \frac{e^{(x-1)^2} - 1}{(x-1)^2} \text{ for } x \neq 1 \text{ and } f(1) = 1. \text{ The function } f \text{ has derivatives of all orders at } x = 1.$$

- (a) Write the first four nonzero terms and the general term of the Taylor series for  $e^{(x-1)^2}$  about  $x = 1$ .
- (b) Use the Taylor series found in part (a) to write the first four nonzero terms and the general term of the Taylor series for  $f$  about  $x = 1$ .
- (c) Use the ratio test to find the interval of convergence for the Taylor series found in part (b).
- (d) Use the Taylor series for  $f$  about  $x = 1$  to determine whether the graph of  $f$  has any points of inflection.
- 

**WRITE ALL WORK IN THE PINK EXAM BOOKLET.**

**END OF EXAM**

更多AP资料下载：<http://www.manfen.net/thread-9638-1-1.html>